






The information in this label chart is no substitute for professional knowledge or judgement and users should exercise their own skill and care with respect to its use. Users are encouraged to consult with other sources, such as the Australian Pharmaceutical Formulary and Handbook and the manufacturer's product information to ensure the information provided in this catalogue chart is suitable for their purpose.

CAUTIONARY & ADVISORY LABELS



WARN 1 1000 per box	WARN 1a 1000 per box	WARN 2 1000 per box	WARN 3a 1000 per box	WARN 3b 1000 per box
 This medicine may cause DROWSINESS and may increase the effects of alcohol. If affected, do not drive a motor vehicle or operate machinery.	 This preparation is to aid sleep. Drowsiness may continue the following day. If affected, do not drive or operate machinery. Avoid alcohol.	 DO NOT TAKE ALCOHOL while being treated with this medicine.	 Take on an empty stomach at least half an hour before meals and at bedtime.	 Take on an empty stomach at least half an hour before food or two hours after food.
WARN 4b 1000 per box	WARN 5 1000 per box	WARN 6 1000 per box	WARN 7a 1000 per box	WARN 7b 1000 per box
 Ask your doctor or pharmacist before taking medicines for heartburn, reflux or indigestion.	 Ask your doctor or pharmacist before using any other medicine including over-the-counter medicines or health products.	 REFRIGERATE Do not freeze	 Discard contents after / /	 Discard days after opening. Date opened / /
WARN 9 1000 per box	WARN 10a 1000 per box	WARN 10b 1000 per box	WARN 11 1000 per box	WARN 12 1000 per box
 RINSE MOUTH with water after each use.	 This medicine replaces Do not use both.	 Active ingredient:  DO NOT USE BOTH	 This medicine may cause dizziness especially when you stand up quickly. Ask your doctor or pharmacist for advice.	 Certain foods and fruit juices should be avoided while being treated with this medicine
WARN 14 1000 per box	WARN 15a 1000 per box	WARN 15b 1000 per box	WARN 16 1000 per box	WARN 18 1000 per box
 Consult your doctor or pharmacist before taking other medicines for pain or inflammation.	 Take once weekly on the same day.	 Special handling and disposal required — ask your pharmacist.	 Use only with approved or recommended device.	 This product has been compounded by the pharmacist.
WARN A 1000 per box	WARN B 1000 per box	WARN C 1000 per box	WARN E 1000 per box	WARN F 1000 per box
 SWALLOW WHOLE Do not crush or chew	 TAKE WITH OR SOON AFTER FOOD	 TAKE AT LEAST HALF AN HOUR BEFORE FOOD	 CONTINUE FOR 14 DAYS AFTER SYMPTOMS CEASE	 TAKE IMMEDIATELY BEFORE FOOD
WARN H 1000 per box	WARN J 1000 per box	WARN K 1000 per box	WARN L 1000 per box	WARN M 1000 per box
 STORE FROZEN	 SHAKE WELL BEFORE EACH USE	 FOR EXTERNAL USE ONLY	 CAUTION NOT TO BE TAKEN	 ALLOW THIS MEDICINE TO DISSOLVE IN YOUR MOUTH
WARN N 1000 per box				


ADDITIONAL ADVISORY LABELS

WARN 61 1000 per box	WARN 62 1000 per box	WARN 62a 1000 per box	WARN 67 1000 per box	WARN 70 1000 per box
 KEEP OUT OF THE REACH OF CHILDREN	 LAST REPEAT FOR FURTHER REPEATS A NEW PRESCRIPTION IS NECESSARY	 LAST REPEAT	 DELIVER	 The Mixture SHAKE THE BOTTLE
WARN 72 1000 per box	WARN 86 1000 per box	WARN 88 1000 per box	WARN 89 1000 per box	WARN 90 1000 per box
 CAUTION NOT TO BE SWALLOWED	 DO NOT TAKE THIS MEDICINE IF ALLERGIC TO PENICILLIN	 TO BE TAKEN UNTIL ALL USED	 PHARMACIST WISHES TO SPEAK TO YOU	 SHAKE WELL
WARN 93 1000 per box	WARN 95 1000 per box	WARN 96 1000 per box	WARN 98 1000 per box	WARN 99 1000 per box
 A LOWER PRICED BRAND OF YOUR MEDICINE IS AVAILABLE PLEASE CONSULT YOUR DOCTOR OR PHARMACIST	 PLEASE GIVE 48 HOURS NOTICE BEFORE NEXT SUPPLY.	 DO NOT USE IF PREGNANT	 PRESCRIPTION OWING FOR THIS ITEM	 THIS MEDICINE MAY DECREASE THE EFFECTIVENESS OF ORAL CONTRACEPTIVES. CONSULT YOUR DOCTOR OR PHARMACIST.
WARN 100 1000 per box	WARN 102 1000 per box	WARN 104 1000 per box	WARN 108 1000 per box	WARN 112 1000 per box
 THIS PREPARATION IS AN ANTIBIOTIC. THE ENTIRE COURSE SHOULD BE COMPLETED.	 SHAKE WELL AND KEEP IN REFRIGERATOR	 INHALE SLOW and STEADY	 INHALE QUICK and DEEP	 Please return empty inhaler to pharmacy for safe disposal
WARN 126 1000 per box	WARN 142 1000 per box	WARN 144 1000 per box	WARN 200 500 per box	WARN 202 500 per box
 A list of inactive ingredients is available from your pharmacist upon request	 Risk of serious harm or death with intravenous administration. Never to be handled by patients. Access restricted to pharmacists and clinicians only.	 DO NOT inject may cause death or serious injury if injected or taken by another person	 POISON! Not to be taken	 FOR EXTERNAL USE ONLY
WARN 140 1000 per box	WARN 205 500 per box	WARN 206 500 per box	WARN 201 500 per box	WARN 203 500 per box

BULK BUY – 12 ASSORTED BOXES AND SAVE

ALTERNATIVE MEDICINE LABELS

CAN301 1000 per box	CAN301A 1000 per box	CAN302 1000 per box	CAN303 1000 per box
 Store in a cool dry place out of sunlight.	 DO NOT VAPE. To be taken orally or sublingually (under the tongue).	 Take doses 6 to 8 hours apart	 May take 1 to 1.5 hrs to take effect
CAN304 1000 per box	CAN305 1000 per box	CAN306 1000 per box	CAN307 1000 per box
 May take 2 to 3 mins to take effect	 DO NOT VAPE. Inhalation of this preparation may be harmful.	 LAST REPEAT	 DO NOT REFRIGERATE
CAN308 1000 per box	CAN309 1000 per box	CAN311 1000 per box	CAN312 1000 per box
 USE WITH A TGA APPROVED VAPORIZING DEVICE	 DAY TIME USE	 NIGHT TIME USE	 Dose Taken Initial Dose Remaining
CAN313 1000 per box	CAN320 1000 per box	CAN321 1000 per box	CAN325 1000 per box



TO PLACE YOUR ORDER CALL **1300 651 118**

E| sales@stirlingfildes.com.au W| www.stirlingfildes.com.au